

**GEOMETRY AND TOPOLOGY PRELIMINARY EXAM**  
**FALL 2020**

[1] Let  $M_2(\mathbb{R})$  be the set of all real  $2 \times 2$  matrices. Consider the subset given by  $SL_2(\mathbb{R}) = \{A \in M_2(\mathbb{R}) \mid \det A = 1\}$ .

(1) Show that  $SL_2(\mathbb{R})$  is a 3-dimensional smooth manifold.

(2) Regarding the tangent space  $T_A SL_2(\mathbb{R})$  at  $A$  as a linear subspace of  $T_A M_2(\mathbb{R}) \cong M_2(\mathbb{R})$ , show that

$$T_A SL_2(\mathbb{R}) = A \cdot \left\{ \begin{pmatrix} p & q \\ r & s \end{pmatrix} \mid p + s = 0 \right\} \subset M_2(\mathbb{R}).$$

[2] On  $\mathbb{R}^3$ , consider a distribution generated by the following vector fields:

$$X = \frac{\partial}{\partial x} + 2x \frac{\partial}{\partial z}, \quad Y = \frac{\partial}{\partial y} + 2y \frac{\partial}{\partial z}, \quad Z = -y \frac{\partial}{\partial x} + 2x \frac{\partial}{\partial y}.$$

(1) Show that the above distribution is an involutive 2-dimensional distribution.

(2) Calculate the integral manifold through the origin.

[3] Let  $\alpha$  and  $\beta$  be differential forms on a manifold  $M$  and let  $X$  be a vector field. Prove that

$$L_X(\alpha \wedge \beta) = (L_X \alpha) \wedge \beta + \alpha \wedge (L_X \beta)$$

where  $L_X$  denotes the Lie derivative.

[4] Is there an onto submersion  $S^3 \rightarrow S^2$ ?

[5] Given a Riemannian manifold  $(M, g)$ , show that for each  $p \in M$ , there exists a neighborhood  $U$  of  $p$  such that every geodesic line in  $U$  is a length minimizing curve.

[6] Let  $S$  be a compact orientable surface of genus 2. Compute the fundamental group  $\pi_1(S)$ . (Hint: What is the fundamental group of a punctured torus?)

[7] Let  $n \geq 0$ . Prove that every continuous map  $f : D^n \rightarrow D^n$  from the closed  $n$ -disk to itself has a fixed point.