

## Geometry and Topology Prelim, UCSC, Fall 2016

1. Consider the vector fields

$$v = y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y} - z^m \frac{\partial}{\partial z}$$

and

$$w = \frac{\partial}{\partial x} + z \frac{\partial}{\partial z}.$$

on  $\mathbb{R}^3$ , where  $m \geq 0$ .

- Find the (local) flow of  $v$ . Is  $v$  complete? (Note: Pay attention to  $m$ .)
- Find the bracket  $[v, w]$ .
- Is the distribution spanned by  $v$  and  $w$  (on the open subset where these vector fields are linearly independent) integrable?

2. Let  $\alpha$  and  $\beta$  be differential forms on a manifold  $M$  and let  $X$  be a vector field. Prove that

$$L_X(\alpha \wedge \beta) = (L_X\alpha) \wedge \beta + \alpha \wedge (L_X\beta).$$

3. Is there an embedding of the Klein bottle in  $\mathbb{R}^4$ ? Construct such an embedding or prove that it does not exist.

4. Consider the special orthogonal group  $\text{SO}(n)$ , i.e.,  $\text{SO}(n)$  is formed by  $n \times n$  matrices  $A$  with  $AA^T = I$  and  $\det A = 1$ .

- Show that  $\text{SO}(n)$  is a smooth manifold.
- Show that  $\text{SO}(n)$  is compact.
- Is the tangent bundle  $T\text{SO}(n)$  trivial? Justify your answer.

5. Let  $(G, *)$  be a topological group with multiplication  $*$ :  $G \times G \rightarrow G$ . In  $G$ , there are two ways to multiply loops based at the identity element  $e \in G$ :

(i) Use the usual concatenation of loops,

$$(f \cdot g)(s) = \begin{cases} f(2s), & 0 \leq s \leq 1/2, \\ g(2s - 1), & 1/2 \leq s \leq 1. \end{cases}$$

(ii) Use the group structure in  $G$ :

$$(f * g)(s) = f(s) * g(s), \quad 0 \leq s \leq 1.$$

Show that  $f \cdot g \simeq f * g$  for all loops  $f$  and  $g$  based at  $e$ . Thus, two ways of multiplying loops define the same product structure in  $\pi_1(G, e)$ . Next show that  $\pi_1(G, e)$  is in fact abelian regardless of whether  $G$  is abelian or not.

Over please!

6. Let  $X$  be the topological space obtained by attaching the boundary  $\partial D$  of the closed unit disc  $D$  to the unit circle via a map  $\varphi: \partial D = S^1 \rightarrow S^1$  of degree three. (For instance, we can take  $\varphi(z) = z^3$ , where we view  $S^1$  as the unit circle in  $\mathbb{C}$ .) Find the homology of  $X$  with integer coefficients. What changes if we use rational coefficients?

7. Consider the upper half-plane  $M = \{(x, y) \in \mathbb{R}^2 \mid y > 0\}$  equipped with the Riemannian metric

$$g = \frac{dx \otimes dx + dy \otimes dy}{y^2}.$$

- (a) Find the equation  $t \mapsto (x(t), y(t))$  of the geodesic starting at  $(0, 1)$  with initial velocity  $(0, 1)$ . (Hints: You can use the fact that the Christoffel symbols of  $g$  are zero except for  $\Gamma_{xx}^y = 1/y$  and  $\Gamma_{xy}^x = \Gamma_{yx}^y = \Gamma_{yy}^y = -1/y$ . You don't need to calculate them. Alternatively, and this is a better method, you can base your argument on the observation that the metric is invariant with respect to the reflection in the  $y$ -axis.)
- (b) What is the distance between the points  $(0, a)$  and  $(0, b)$ ?